

# Ferro

Breed: Lagotto Romagnolo  
Microchip number: 978101084931433  
Birth date: 2024-07-26

Registration number: FI39404/24  
Test date: 2024-12-14  
ID kit: DKGQLJC



## Ferro's Profile

### Pet information

Registered name	Sex
Lagonotella Il Tesoro del Mare	M
Owner reported breed	Date of birth
Lagotto Romagnolo	2024-07-26
Microchip number	
978101084931433	

### Genetic Diversity

Ferro's Percentage of Heterozygosity  
39%

### Health summary

- At Risk 0 conditions
- Carrier 0 conditions
- Clear 272 conditions



# Ferro

Breed: Lagotto Romagnolo  
Microchip number: 978101084931433  
Birth date: 2024-07-26

Registration number: FI39404/24  
Test date: 2024-12-14  
ID kit: DKGQLJC



## Genetic Diversity

### Heterozygosity

#### Ferro's Percentage of Heterozygosity

39%

Ferro's genome analysis shows higher than average genetic heterozygosity when compared with other Lagotto Romagnolo - Romagna Water Dogs.

#### Typical Range for Lagotto Romagnolo - Romagna Water Dogs

32% - 38%





## Health conditions known in the breed

Benign Familial Juvenile Epilepsy	Gene	Risk Variant	Copies	Inheritance	Result
	LGI2	A>T	0	AR	Clear

Information about the genetic condition

Onset of epileptic seizures in BFJE is between 5 to 9 weeks of age. The seizures consist of whole-body tremors, ataxia (uncoordinated movements), and stiffness. Some affected puppies are able to stand and/or walk during a seizure while others are recumbent and unable to stand. The epileptic signs can sometimes be associated with alterations of consciousness. Frequency of seizures can vary between individuals: there can be several seizures a day or sporadic seizures a few times a week. Usually an affected puppy seems completely normal between the seizures. However, in severe cases there can be neurological signs, such as generalized ataxia between the seizures. The BFJE form of epilepsy is relatively benign since the seizures typically end by 4 months of age.

Breeder recommendation

This disease is autosomal recessive meaning that two copies of the mutation are needed for disease signs to develop. However dogs with one copy of the disease mutation have been reported as having a slightly increased chance of developing this disease. Breeding a carrier dog with one copy of the BFJE mutation with a clear dog with no copies of the BFJE mutation, will produce a litter with around half carrier puppies and half clear puppies. Puppies in a litter which is expected to contain carriers should be tested prior to breeding. Carrier to carrier matings are not advised as the resulting litter may contain affected puppies. Please note: It is possible that disease signs similar to the ones caused by the BFJE mutation could develop due to a different genetic or clinical cause.

Hyperuricosuria	Gene	Risk Variant	Copies	Inheritance	Result
	SLC2A9	G>T	0	AR	Clear

Information about the genetic condition

HUU predisposes affected dogs to the formation of urate stones. Clinical signs of urolithiasis include hematuria, pain while urinating, and blockage of the urinary tract. Patients with urinary stones are more susceptible to urinary tract infections. Blockage of the urinary tract is a life-threatening condition that requires immediate veterinary care. In Dalmatians, the clinical signs are more common in males than in females. As many as 34% of all male Dalmatians are diagnosed with urate stones.

Breeder recommendation

This disease is autosomal recessive meaning that two copies of the mutation are needed for disease signs to occur. A carrier dog with one copy of the HUU mutation can be safely bred with a clear dog with no copies of the HUU mutation. About half of the puppies will have one copy (carriers) and half will have no copies of the HUU mutation. A dog with two copies of the HUU mutation can be safely bred with a clear dog. The resulting puppies will all be carriers. Puppies in a litter which is expected to contain carriers should be tested prior to breeding. In some breeds, such as the Dalmatian, the frequency of the disease mutation is very high. Carriers and dogs with two copies of the disease mutation (genetically affected dogs) should be used for breeding purposes, with the aim of gradually reducing the frequency of the mutant gene within the breed population. Where possible, matings should be avoided that would result in litters that could contain dogs with two copies of the disease mutation, such as a mating between two dogs with two copies of the HUU mutation or between a dog with one copy and a dog with two copies of the HUU mutation. Please note: It is possible that disease signs similar to the ones caused by the HUU mutation could develop due to a different genetic or clinical cause.



Breed: Lagotto Romagnolo  
Microchip number: 978101084931433  
Birth date: 2024-07-26

Registration number: FI39404/24  
Test date: 2024-12-14  
ID kit: DKGQLJC



## Health conditions known in the breed

Lagotto Storage Disease	Gene	Risk Variant	Copies	Inheritance	Result
	ATG4D	G>A	0	AR	Clear

### Information about the genetic condition

The onset of clinical signs and pattern of progression vary considerably between individuals. The first signs can be seen at the age of 4 months to 4 years. Typically the first observable sign of LSD is usually progressive ataxia (uncoordinated movements), which may not be noticed by the owner. Some of the affected dogs also suffer from episodic nystagmus (involuntary eye movement). In some cases, nystagmus is the first sign noticed. Lagotto storage disease is a progressive condition leading to behavioral changes such as restlessness, depression, and aggression. The life expectancy of affected dogs depends on the progression of the disorder and the severity of the signs. Some dogs can live for several years with mild signs, but dogs with severe clinical signs are usually euthanized earlier.

### Breeder recommendation

This disease is autosomal recessive meaning that two copies of the mutation are needed for disease signs to occur. A carrier dog with one copy of the LSD mutation can be safely bred with a clear dog with no copies of the LSD mutation. About half of the puppies will have one copy (carriers) and half will have no copies of the LSD mutation. Puppies in a litter which is expected to contain carriers should be tested prior to breeding. Carrier to carrier matings are not advised as the resulting litter may contain affected puppies. Please note: It is possible that disease signs similar to the ones caused by the LSD mutation could develop due to a different genetic or clinical cause.

Progressive Rod Cone Degeneration (prcd-PRA)	Gene	Risk Variant	Copies	Inheritance	Result
	PRCD	G>A	0	AR	Clear

### Information about the genetic condition

Clinical signs of PRCD are related to progressive loss of function of rod photoreceptors, followed by loss of function of cone photoreceptors. Typical signs of disease include hyper-reflective tapetum and attenuated blood vessels. Age of onset for this form of PRA is generally early adulthood, although exact age of onset may vary significantly among different breeds. The disorder is progressive, causing increasing levels of vision loss and eventual blindness.

### Breeder recommendation

This disease is autosomal recessive meaning that two copies of the mutation are needed for disease signs to occur. A carrier dog with one copy of the prcd-PRA mutation can be safely bred with a clear dog with no copies of the prcd-PRA mutation. About half of the puppies will have one copy (carriers) and half will have no copies of the prcd-PRA mutation. A dog with two copies of the prcd-PRA mutation can be safely bred with a clear dog. The resulting puppies will all be carriers. Puppies in a litter which is expected to contain carriers should be tested prior to breeding. Carrier to carrier matings are not advised as the resulting litter may contain affected puppies. Please note: It is possible that disease signs similar to the ones caused by the prcd-PRA mutation could develop due to a different genetic or clinical cause.





Traits

Coat Color

	Gene	Variant	Copies	Result
<b>Fawn</b>	ASIP	ay	0	No effect
<b>Recessive Black</b>	ASIP	a	0	No effect
<b>Tan Points</b> Two copies, or occasionally one copy, of this variant may result in a black and tan coat color pattern.	ASIP	at	2	Tan points possible
<b>Dominant Black</b> One or two copies of the dominant black will give a dog a black coat (depending on other variants), black eye rims, nose and pads. One copy may also give a tiger striped appearance, known as brindle patterning.	CBD103	KB	2	Black possible
<b>Mask</b>	MC1R	Em	0	No effect
<b>Recessive Red (e1)</b> To show a solid red coat, a dog must inherit two copies of a Recessive Red variant, one from each parent. This can either be two copies of a particular variant, such as this one (e1) or two of any combination of recessive red variants. Recessive red coats will appear white, cream, yellow or red, although there are other variants that can result in a similar appearance. The amount of red pigment in the coat, called the intensity, is governed by other genes.	MC1R	e1	2	Cream to red coat likely
<b>Recessive Red (e2)</b>	MC1R	e2	0	No effect
<b>Recessive Red (e3)</b>	MC1R	e3	0	No effect
<b>Sable (Discovered in the Cocker Spaniel)</b>	MC1R	eH	0	No effect
<b>Widow's Peak (Discovered in Ancient dogs)</b>	MC1R	eA	0	No effect
<b>Widow's Peak (Discovered in the Afghan Hound and Saluki)</b>	MC1R	EG	0	No effect

Color Modification

	Gene	Variant	Copies	Result
<b>Cocoa (Discovered in the French Bulldog)</b>	HPS3	co	0	No effect





Color Modification

	Gene	Variant	Copies	Result
<b>Red Intensity</b>  Dogs with two copies of the Red Intensity variant are more likely to show yellow, cream or white coat shades instead of deeper red shades. If the dog does not display solid red or red coat patterns, there will be no visible effect. Other genes, notably variants in the KITLG gene, are also thought to contribute to red pigment intensity variation, so some dogs may have yellow or buff colored coats.	MFSD12	i	2	White to yellow coat shades likely
<b>Dilution (d1) Linkage test</b>	MLPH	d <sup>1</sup>	0	No effect
<b>Dilution (d2)</b>	MLPH	d <sup>2</sup>	0	No effect
<b>Dilution (d3)</b>	MLPH	d <sup>3</sup>	0	No effect
<b>Chocolate (basd)</b>	TYRP1	b <sup>asd</sup>	0	No effect
<b>Chocolate (bc)</b>  To show chocolate coloration a dog must inherit two chocolate variants, one from each parent. This can either be two copies of a particular variant, such as this one ("bc"), or two of any combination of chocolate variants.	TYRP1	b <sup>c</sup>	1	Black features likely, chocolate possible
<b>Chocolate (bd)</b>  To show chocolate coloration a dog must inherit two chocolate variants, one from each parent. This can either be two copies of a particular variant, such as this one ("bd"), or two of any combination of chocolate variants. This variant is unique in that it can occur on the same chromosome as another chocolate variant, where both variants are donated from one parent. If the other parent does not also donate a chocolate variant, the dog will still express black pigment, not chocolate.	TYRP1	b <sup>d</sup>	1	Black features likely, chocolate possible
<b>Chocolate (be)</b>	TYRP1	b <sup>e</sup>	0	No effect
<b>Chocolate (bh)</b>	TYRP1	b <sup>h</sup>	0	No effect
<b>Chocolate (bs)</b>  To show chocolate coloration a dog must inherit two chocolate variants, one from each parent. This can either be two copies of a particular variant, such as this one ("bs"), or two of any combination of chocolate variants.	TYRP1	b <sup>s</sup>	1	Black features likely, chocolate possible



Breed: Lagotto Romagnolo  
Microchip number: 978101084931433  
Birth date: 2024-07-26

Registration number: FI39404/24  
Test date: 2024-12-14  
ID kit: DKGQLJC



Coat Patterns

	Gene	Variant	Copies	Result
<b>Piebald</b> Dog with copies of the Piebald variant are likely to show white spotting, patches and/or a white coat, with two copies having a greater effect than one, although the strength of this effect may be influenced by other genes.	MITF	S <sup>p</sup>	2	Particolor or white markings possible
<b>Merle</b>	PMEL	M	0	No effect
<b>Harlequin</b>	PSMB7	H	0	No effect
<b>Saddle Tan</b>	RALY	-	0	No effect
<b>Roan (Linkage test)</b>	USH2A	Tr	0	No effect

Coat Length and Curl

	Gene	Variant	Copies	Result
<b>Long Hair (lh1)</b> To show a long coat, a dog must inherit two copies of a Long Hair variant, one from each parent. This can either be two copies of a particular variant, such as this one (lh1) or two of any combination of long hair variants. However, there are other variants suspected to influence coat length.	FGF5	lh <sup>1</sup>	2	Long coat
<b>Long Hair (lh2)</b>	FGF5	lh <sup>2</sup>	0	No effect
<b>Long Hair (lh3)</b>	FGF5	lh <sup>3</sup>	0	No effect
<b>Long Hair (lh4)</b>	FGF5	lh <sup>4</sup>	0	No effect
<b>Long Hair (lh5)</b>	FGF5	lh <sup>5</sup>	0	No effect
<b>Curly Coat</b> One copy of this variant is likely to give a soft curl or wave whereas two copies are likely to give a tighter curl. A curly coat is less apparent in dogs with short hair than those with long. There is one other known Curl variant, and likely other unknown variants that exist.	KRT71	C	1	Soft curl or wave likely





Hairlessness

	Gene	Variant	Copies	Result
Hairlessness (Discovered in the Chinese Crested Dog) Linkage test	FOXI3	Hr <sup>cc</sup>	0	No effect
Hairlessness (Discovered in the American Hairless Terrier)	SGK3	hra <sup>ht</sup>	0	No effect
Hairlessness (Discovered in the Scottish Deerhound)	SKG3	hr <sup>sd</sup>	0	No effect

Shedding

	Gene	Variant	Copies	Result
Reduced Shedding  One or two copies of the Reduced Shedding variant is likely to reduce a dog's tendency to shed. Copies of the Furnishings variant, particularly two, also reduce the tendency of a dog to shed.	MC5R	sd	1	Occasional shedder

More Coat Traits

	Gene	Variant	Copies	Result
Hair Ridge	FGF3, FGF4, FGF19, ORAOV1	R	0	No effect
Furnishings  Dogs with one or two copies of the Furnishing variant are likely to display a fuzzy beard, moustache and eyebrows, but a long or curly coat will make this variant less apparent.	RSPO2	F	2	Furnishings likely
Albino	SLC45A2	cal	0	No effect

Head Shape

	Gene	Variant	Copies	Result
Short Snout (BMP3 variant)	BMP3	-	0	No effect
Short Snout (SMOC2 variant)	SMOC2	-	0	No effect





Eye Color

	Gene	Variant	Copies	Result
Blue Eyes (Discovered in the Siberian Husky)	ALX4	-	0	No effect

Ears

	Gene	Variant	Copies	Result
<b>Floppy Ears</b> Dogs with zero copies of this variant are more likely to have permanently upright or prick ears, and fully folded ears are more likely with two copies inherited. Please note however that many genetic variants influence ear carriage. Dogs with some cartilage stiffness to their ears can sometimes raise their ears upright when 'at alert' but will flop down when relaxed.	MSRB3	-	2	Floppy ears more likely

Extra Toes

	Gene	Variant	Copies	Result
Hind Dewclaws (Discovered in Asian breeds)	LMBR1	DC-1	0	No effect
Hind Dewclaws (Discovered in Western breeds)	LMBR1	DC-2	0	No effect

More Body Features

	Gene	Variant	Copies	Result
Back Muscle and Bulk	ACSL4	-	0	No effect
High Altitude Adaptation	EPAS1	-	0	No effect
Short Legs (Chondrodysplasia, CDPA)	FGF4	-	0	No effect
Short Legs (Chondrodystrophy, CDDY)	FGF4	-	0	No effect
Short Tail	T-box	T	0	Full tail length likely





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
2,8-dihydroxyadenine (DHA) Urolithiasis	APRT	G>A	0	AR	Clear
Acral Mutilation Syndrome	GDNF	C>T	0	AR	Clear
Acute Respiratory Distress Syndrome	ANLN	C>T	0	AR	Clear
Alaskan Husky Encephalopathy	SLC19A3	G>A	0	AR	Clear
Alexander Disease	GFAP	G>A	0	AR	Clear
Amelogenesis Imperfecta (Discovered in the Italian Greyhound)	ENAM	Deletion	0	AR	Clear
Amelogenesis Imperfecta (Discovered in the Lancashire Heeler)	Confidential	-	0	AR	Clear
Amelogenesis Imperfecta (Discovered in the Parson Russell Terrier)	ENAM	C>T	0	AR	Clear
Bandera's Neonatal Ataxia	GRM1	Insertion	0	AR	Clear
Bernard-Soulier Syndrome (Discovered in the Cocker Spaniel)	GP9	Deletion	0	AR	Clear
Canine Congenital Stationary Night Blindness (Discovered in the Beagle)	LRIT3	Deletion	0	AR	Clear
Canine Leukocyte Adhesion Deficiency (CLAD), type III	FERMT3	Insertion	0	AR	Clear
Canine Multifocal Retinopathy 1	BEST1	C>T	0	AR	Clear
Canine Multifocal Retinopathy 2	BEST1	G>A	0	AR	Clear
Canine Multifocal Retinopathy 3	BEST1	Deletion	0	AR	Clear
Canine Multiple Systems Degeneration (Discovered in the Chinese Crested Dog)	SERAC1	Deletion	0	AR	Clear
Canine Scott Syndrome	ANO6	G>A	0	AR	Clear
Cardiomyopathy and Juvenile Mortality (Discovered in the Belgian Shepherd)	YARS2	G>A	0	AR	Clear
Centronuclear Myopathy (Discovered in the Great Dane)	BIN1	A>G	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Centronuclear Myopathy (Discovered in the Labrador Retriever)	PTPLA	Insertion	0	AR	Clear
Cerebellar Ataxia	RAB24	A>C	0	AR	Clear
Cerebellar Cortical Degeneration	SNX14	C>T	0	AR	Clear
Cerebellar Hypoplasia	VLDLR	Deletion	0	AR	Clear
Cerebral Dysfunction	SLC6A3	G>A	0	AR	Clear
Chondrodysplasia (Discovered in Norwegian Elkhound and Karelian Bear Dog)	ITGA10	C>T	0	AR	Clear
Chondrodystrophy (CDDY) and Intervertebral Disc Disease (IVDD) Risk	FGF4 retrogene	Insertion	0	AD	Clear
Cleft Lip & Palate with Syndactyly	ADAMTS20	Deletion	0	AR	Clear
Cleft Palate	DLX6	C>A	0	AR	Clear
CNS Atrophy with Cerebellar Ataxia (Discovered in the Belgian Shepherd)	SEPP1	Deletion	0	AR	Clear
Coat Color Dilution and Neurological Defects (Discovered in the Miniature Dachshund)	MYO5A	Insertion	0	AR	Clear
Collie Eye Anomaly (CEA)	NHEJ1	Deletion	0	AR	Clear
Complement 3 Deficiency	C3	Deletion	0	AR	Clear
Cone Degeneration (Discovered in the Alaskan Malamute)	CNGB3	Deletion	0	AR	Clear
Cone Degeneration (Discovered in the German Shepherd Dog)	CNGA3	C>T	0	AR	Clear
Cone Degeneration (Discovered in the German Shorthaired Pointer)	CNGB3	G>A	0	AR	Clear
Cone-Rod Dystrophy	NPHP4	Deletion	0	AR	Clear
Cone-Rod Dystrophy 1	PDE6B	Deletion	0	AR	Clear
Cone-Rod Dystrophy 2	IQCB1	Insertion	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Congenital Cornification (Discovered in the Labrador Retriever)	NSDHL	Deletion	0	XD	Clear
Congenital Dys hormonogenic Hypothyroidism with Goiter (Discovered in the Shih Tzu)	SLC5A5	G>A	0	AR	Clear
Congenital Eye Malformations (Discovered in the Golden Retriever)	SIX6	C>T	0	AD	Clear
Congenital Hypothyroidism (Discovered in the Tenterfield Terrier)	TPO	C>T	0	AR	Clear
Congenital Hypothyroidism (Discovered in the Toy Fox and Rat Terrier)	TPO	C>T	0	AR	Clear
Congenital Muscular Dystrophy (Discovered in the Italian Greyhound)	LAMA2	G>A	0	AR	Clear
Congenital Muscular Dystrophy (Discovered in the Staffordshire Bull Terrier)	LAMA2	Deletion	0	AR	Clear
Congenital Myasthenic Syndrome (Discovered in the Golden Retriever)	COLQ	G>A	0	AR	Clear
Congenital Myasthenic Syndrome (Discovered in the Heideterrier)	CHRNE	Insertion	0	AR	Clear
Congenital Myasthenic Syndrome (Discovered in the Jack Russell Terrier)	CHRNE	Insertion	0	AR	Clear
Congenital Myasthenic Syndrome (Discovered in the Labrador Retriever)	COLQ	T>C	0	AR	Clear
Congenital Myasthenic Syndrome (Discovered in the Old Danish Pointer)	CHAT	G>A	0	AR	Clear
Congenital Stationary Night Blindness (CSNB)	RPE65	A>T	0	AR	Clear
Craniomandibular Osteopathy (Discovered in Scottish Terrier breeds)	SLC37A2	C>T	0	AD	Clear
Craniomandibular Osteopathy (Discovered in the Australian Terrier)	COL1A1	C>T	0	AD	Clear
Craniomandibular Osteopathy (Discovered in the Basset Hound)	SLC37A2	C>T	0	AD	Clear
Craniomandibular Osteopathy (Discovered in the Weimaraner)	SLC35D1	Deletion	0	AD	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Cystic Renal Dysplasia and Hepatic Fibrosis	INPP5E	G>A	0	AR	Clear
Cystinuria Type I-A	SLC3A1	C>T	0	AR	Clear
Cystinuria Type II-A	SLC3A1	Deletion	0	AD	Clear
Darier Disease (Discovered in the Irish Terrier)	ATP2A2	Insertion	0	AD	Clear
Deafness and Vestibular Dysfunction (DINGS1), (Discovered in Doberman Pinscher)	PTPRQ	Insertion	0	AR	Clear
Deafness and Vestibular Dysfunction (DINGS2), (Discovered in Doberman Pinscher)	MYO7A	G>A	0	AR	Clear
Degenerative Myelopathy	SOD1	G>A	0	AR	Clear
Demyelinating Neuropathy	SBF2	G>T	0	AR	Clear
Dental Hypomineralization	FAM20C	C>T	0	AR	Clear
Dental-Skeletal-Retinal Anomaly (Discovered in the Cane Corso)	MIA3	Deletion	0	AR	Clear
Dilated Cardiomyopathy (Discovered in the Schnauzer)	RBM20	Deletion	0	AR	Clear
Disproportionate Dwarfism (Discovered in the Dogo Argentino)	PRKG2	C>A	0	AR	Clear
Dominant Progressive Retinal Atrophy	RHO	C>G	0	AD	Clear
Dystrophic Epidermolysis Bullosa (Discovered in the Basset Hound)	COL7A1	Insertion	0	AR	Clear
Dystrophic Epidermolysis Bullosa (Discovered in the Central Asian Ovcharka)	COL7A1	C>T	0	AR	Clear
Dystrophic Epidermolysis Bullosa (Discovered in the Golden Retriever)	COL7A1	C>T	0	AR	Clear
Early Adult Onset Deafness For Border Collies only (Linkage test)	Intergenic	Insertion	0	AR	Clear
Early Retinal Degeneration (Discovered in the Norwegian Elkhound)	STK38L	Insertion	0	AR	Clear
Early-Onset Adult Deafness (Discovered in the Rhodesian Ridgeback)	EPS8L2	Deletion	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Early-Onset Progressive Polyneuropathy (Discovered in the Alaskan Malamute)	NDRG1	G>T	0	AR	Clear
Early-Onset Progressive Polyneuropathy (Discovered in the Greyhound)	NDRG1	Deletion	0	AR	Clear
Early-Onset Progressive Retinal Atrophy (Discovered in the Portuguese Water Dog)	CCDC66	Insertion	0	AR	Clear
Early-Onset Progressive Retinal Atrophy, (Discovered in the Spanish Water Dog)	PDE6B	Deletion	0	AR	Clear
Ehlers-Danlos Syndrome (Discovered in mixed breed)	COL5A1	G>A	0	AD	Clear
Ehlers-Danlos Syndrome (Discovered in the Labrador Retriever)	COL5A1	Deletion	0	AD	Clear
Epidermolytic Hyperkeratosis	KRT10	G>T	0	AR	Clear
Episodic Falling Syndrome	BCAN	Insertion	0	AR	Clear
Exercise-Induced Collapse	DNM1	G>T	0	AR	Clear
Factor VII Deficiency	F7	G>A	0	AR	Clear
Factor XI Deficiency	FXI	Insertion	0	AD	Clear
Familial Nephropathy (Discovered in the English Cocker Spaniel)	COL4A4	A>T	0	AR	Clear
Familial Nephropathy (Discovered in the English Springer Spaniel)	COL4A4	C>T	0	AR	Clear
Fanconi Syndrome	FAN1	Deletion	0	AR	Clear
Fetal Onset Neuroaxonal Dystrophy	MFN2	G>C	0	AR	Clear
Focal Non-Epidermolytic Palmoplantar Keratoderma	KRT16	G>C	0	AR	Clear
Generalized Progressive Retinal Atrophy (Discovered in the Schapendoes)	CCDC66	Insertion	0	AR	Clear
Glanzmann Thrombasthenia Type I (Discovered in Great Pyrenees)	ITGA2B	C>G	0	AR	Clear
Glanzmann Thrombasthenia Type I (Discovered in mixed breed dogs)	ITGA2B	C>T	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Globoid Cell Leukodystrophy (Discovered in Terriers)	GALC	A>C	0	AR	Clear
Globoid Cell Leukodystrophy (Discovered in the Irish Setter)	GALC	A>T	0	AR	Clear
Glycogen Storage Disease Type Ia (Discovered in the German Pinscher)	G6PC	Insertion	0	AR	Clear
Glycogen Storage Disease Type Ia (Discovered in the Maltese)	G6PC	G>C	0	AR	Clear
Glycogen Storage Disease Type IIIa, (GSD IIIa)	AGL	Deletion	0	AR	Clear
GM1 Gangliosidosis (Discovered in the Portuguese Water Dog)	GLB1	G>A	0	AR	Clear
GM1 Gangliosidosis (Discovered in the Shiba)	GLB1	Deletion	0	AR	Clear
GM2 Gangliosidosis (Discovered in the Japanese Chin)	HEXA	G>A	0	AR	Clear
GM2 Gangliosidosis (Discovered in the Toy Poodle)	HEXB	Deletion	0	AR	Clear
Hemophilia A (Discovered in Old English Sheepdog)	FVIII	C>T	0	XR	Clear
Hemophilia A (Discovered in the Boxer)	FVIII	C>G	0	XR	Clear
Hemophilia A (Discovered in the German Shepherd Dog - Variant 1)	FVIII	G>A	0	XR	Clear
Hemophilia A (Discovered in the German Shepherd Dog - Variant 2)	FVIII	G>A	0	XR	Clear
Hemophilia A (Discovered in the Havanese)	FVIII	Insertion	0	XR	Clear
Hemophilia A (Discovered in the Labrador Retriever)	Confidential	-	0	XR	Clear
Hemophilia B	FIX	G>A	0	XR	Clear
Hemophilia B (Discovered in the Airedale Terrier)	FIX	Insertion	0	XR	Clear
Hemophilia B (Discovered in the Lhasa Apso)	FIX	Deletion	0	XR	Clear
Hereditary Ataxia (Discovered in the Belgian Malinois)	SLC12A6	Insertion	0	AR	Clear
Hereditary Ataxia (Discovered in the Norwegian Buhund)	KCNIP4	T>C	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Hereditary Calcium Oxalate Urolithiasis, Type 1	Confidential	-	0	AR	Clear
Hereditary Elliptocytosis	SPTB	C>T	0	AD	Clear
Hereditary Footpad Hyperkeratosis	FAM83G	G>C	0	AR	Clear
Hereditary Nasal Parakeratosis (Discovered in the Greyhound)	SUV39H2	Deletion	0	AR	Clear
Hereditary Nasal Parakeratosis (Discovered in the Labrador Retriever)	SUV39H2	A>C	0	AR	Clear
Hereditary Vitamin D-Resistant Rickets Type II	VDR	Deletion	0	AR	Clear
Hypocatalasia	CAT	G>A	0	AR	Clear
Hypomyelination	FNIP2	Deletion	0	AR	Clear
Hypophosphatasia	Confidential	-	0	AR	Clear
Ichthyosis (Discovered in the American Bulldog)	NIPAL4	Deletion	0	AR	Clear
Ichthyosis (Discovered in the Great Dane)	SLC27A4	G>A	0	AR	Clear
Ichthyosis Type 2 (Discovered in the Golden Retriever)	ABHD5	Deletion	0	AR	Clear
Inflammatory Myopathy (Discovered in the Dutch Shepherd Dog)	SLC25A12	A>G	0	AR	Clear
Inflammatory Pulmonary Disease (Discovered in the Rough Collie)	AKNA	Deletion	0	AR	Clear
Intestinal Cobalamin Malabsorption (Discovered in the Beagle)	CUBN	Deletion	0	AR	Clear
Intestinal Cobalamin Malabsorption (Discovered in the Border Collie)	CUBN	Deletion	0	AR	Clear
Intestinal Cobalamin Malabsorption (Discovered in the Komondor)	CUBN	G>A	0	AR	Clear
Intestinal Lipid Malabsorption (Discovered in the Australian Kelpie)	ACSL5	Deletion	0	AR	Clear
Junctional Epidermolysis Bullosa (Discovered in the Australian Cattle Dog Mix)	LAMA3	T>A	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Junctional Epidermolysis Bullosa (Discovered in the Australian Shepherd)	LAMB3	A>G	0	AR	Clear
Juvenile Cataract (Discovered in the Wirehaired Pointing Griffon)	FYCO1	Deletion	0	AR	Clear
Juvenile Dilated Cardiomyopathy (Discovered in the Toy Manchester Terrier)	ABCC9	G>A	0	AR	Clear
Juvenile Encephalopathy (Discovered in the Parson Russell Terrier)	Confidential	-	0	AR	Clear
Juvenile Laryngeal Paralysis and Polyneuropathy	RAB3GAP1	Deletion	0	AR	Clear
Juvenile Myoclonic Epilepsy	DIRAS1	Deletion	0	AR	Clear
L-2-Hydroxyglutaric aciduria (Discovered in the Staffordshire Bull Terrier)	L2HGDH	T>C	0	AR	Clear
L-2-Hydroxyglutaric Aciduria (Discovered in the West Highland White Terrier)	Confidential	-	0	AR	Clear
Lafora Disease (Linkage test)	NHLRC1	Insertion	0	AR	Clear
Lamellar Ichthyosis	TGM1	Insertion	0	AR	Clear
Laryngeal Paralysis (Discovered in the Bull Terrier and Miniature Bull Terrier)	RAPGEF6	Insertion	0	AR	Clear
Leigh-like Subacute Necrotizing Encephalopathy (Discovered in the Yorkshire Terrier)	SLC19A3	Insertion	0	AR	Clear
Lethal Acrodermatitis (Discovered in the Bull Terrier)	MKLN1	A>C	0	AR	Clear
Leukodystrophy (Discovered in the Standard Schnauzer)	TSEN54	C>T	0	AR	Clear
Ligneous Membranitis	PLG	T>A	0	AR	Clear
Limb-girdle Muscular Dystrophy (Discovered in the Boston Terrier)	SGCD	Deletion	0	AR	Clear
Limb-girdle Muscular Dystrophy, Type L3 (Discovered in the Miniature Dachshund)	SGCA	G>A	0	AR	Clear
Lung Developmental Disease (Discovered in the Airedale Terrier)	LAMP3	C>T	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Macrothrombocytopenia (Discovered in Norfolk and Cairn Terrier)	TUBB1	G>A	0	AR	Clear
May-Hegglin Anomaly	MYH9	G>A	0	AD	Clear
MDR1 Medication Sensitivity	MDR1/ABCB1	Deletion	0	AD	Clear
Microphthalmia (Discovered in the Soft-Coated Wheaten Terrier)	RBP4	Deletion	0	AR	Clear
Mucopolysaccharidosis Type IIIA (Discovered in the Dachshund)	SGSH	C>A	0	AR	Clear
Mucopolysaccharidosis Type IIIA (Discovered in the New Zealand Huntaway)	SGSH	Insertion	0	AR	Clear
Mucopolysaccharidosis Type VII (Discovered in the Brazilian Terrier)	GUSB	C>T	0	AR	Clear
Mucopolysaccharidosis Type VII (Discovered in the German Shepherd Dog)	GUSB	G>A	0	AR	Clear
Mucopolysaccharidosis VI (Discovered in the Miniature Pinscher)	ARSB	G>A	0	AR	Clear
Muscular Dystrophy (Discovered in the Cavalier King Charles Spaniel)	Dystrophin	G>T	0	XR	Clear
Muscular Dystrophy (Discovered in the Golden Retriever)	Dystrophin	A>G	0	XR	Clear
Muscular Dystrophy (Discovered in the Landseer)	COL6A1	G>T	0	AR	Clear
Muscular Dystrophy (Discovered in the Norfolk Terrier)	Dystrophin	Deletion	0	XR	Clear
Muscular Dystrophy-Dystroglycanopathy (Discovered in the Labrador Retriever)	LARGE	C>T	0	AR	Clear
Muscular Hypertrophy (Double Muscling)	MSTN	T>A	0	AR	Clear
Musladin-Lueke Syndrome	ADAMTSL2	C>T	0	AR	Clear
Myeloperoxidase Deficiency	MOP	C>T	0	AR	Clear
Myotonia Congenita (Discovered in Australian Cattle Dog)	CLCN1	Insertion	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Myotonia Congenita (Discovered in the Labrador Retriever)	CLCN1	T>A	0	AR	Clear
Myotonia Congenita (Discovered in the Miniature Schnauzer)	CLCN1	C>T	0	AR	Clear
Myotubular Myopathy	MTM1	A>C	0	XR	Clear
Narcolepsy (Discovered in the Dachshund)	HCRT2	G>A	0	AR	Clear
Narcolepsy (Discovered in the Labrador Retriever)	HCRT2	G>A	0	AR	Clear
Nemaline Myopathy	NEB	C>A	0	AR	Clear
Neonatal Cerebellar Cortical Degeneration	SPTBN2	Deletion	0	AR	Clear
Neonatal Encephalopathy with Seizures	ATF2	T>G	0	AR	Clear
Neuroaxonal Dystrophy (Discovered in Spanish Water Dog)	TECPR2	C>T	0	AR	Clear
Neuroaxonal Dystrophy (Discovered in the Papillon)	PLA2G6	G>A	0	AR	Clear
Neuroaxonal Dystrophy (Discovered in the Rottweiler)	VPS11	A>G	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 1	PPT1	Insertion	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 12 (Discovered in the Australian Cattle Dog)	ATP13A2	C>T	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 5 (Discovered in the Border Collie)	CLN5	C>T	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 5 (Discovered in the Golden Retriever)	CLN5	Deletion	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 7	MFSD8	Deletion	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 8 (Discovered in the Alpine Dachsbracke)	CLN8	Deletion	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 8 (Discovered in the Australian Shepherd)	CLN8	G>A	0	AR	Clear
Neuronal Ceroid Lipofuscinosis 8 (Discovered in the English Setter)	CLN8	T>C	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Neuronal Ceroid Lipofuscinosis 8 (Discovered in the Saluki)	CLN8	Insertion	0	AR	Clear
Obesity risk (POMC)	POMC	Deletion	0	AD	Clear
Osteochondrodysplasia	SLC13A1	Deletion	0	AR	Clear
Osteochondromatosis (Discovered in the American Staffordshire Terrier)	EXT2	C>A	0	AR	Clear
Osteogenesis Imperfecta (Discovered in the Beagle)	COL1A2	C>T	0	AD	Clear
Osteogenesis Imperfecta (Discovered in the Dachshund)	SERPINH1	T>C	0	AR	Clear
P2RY12-associated Bleeding Disorder	P2RY12	Deletion	0	AR	Clear
Palmoplantar Hyperkeratosis (Discovered in the Rottweiler)	DSG1	Deletion	0	AR	Clear
Paroxysmal Dyskinesia	PIGN	C>T	0	AR	Clear
Persistent Müllerian Duct Syndrome	AMHR2	C>T	0	AR	Clear
Phosphofructokinase Deficiency	PFKM	G>A	0	AR	Clear
Pituitary Dwarfism (Discovered in the Karelian Bear Dog)	POU1F1	C>A	0	AR	Clear
Polycystic Kidney Disease	PKD1	G>A	0	AD	Clear
Prekallikrein Deficiency	KLKB1	T>A	0	AR	Clear
Primary Ciliary Dyskinesia	CCDC39	C>T	0	AR	Clear
Primary Ciliary Dyskinesia (Discovered in the Alaskan Malamute)	NME5	Deletion	0	AR	Clear
Primary Lens Luxation	ADAMTS17	G>A	0	AR	Clear
Primary Open Angle Glaucoma (Discovered in Basset Fauve de Bretagne)	ADAMTS17	G>A	0	AR	Clear
Primary Open Angle Glaucoma (Discovered in Petit Basset Griffon Vendeen)	ADAMTS17	Insertion	0	AR	Clear
Primary Open Angle Glaucoma and Lens Luxation (Discovered in Chinese Shar-Pei)	ADAMTS17	Deletion	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Progressive Early-Onset Cerebellar Ataxia	SEL1L	T>C	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Basenji)	SAG	T>C	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Golden Retriever - GR-PRA 2 variant)	TTC8	Deletion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Golden Retriever - GR-PRA1 variant)	SLC4A3	Insertion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Lapponian Herder)	IFT122	C>T	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Lhasa Apso)	IMPG2	Insertion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Miniature Long Haired Dachshund)	RPGRIP1	Insertion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Papillon and Phalène)	CNGB1	Deletion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Shetland Sheepdog - BBS2 variant)	Confidential	-	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Shetland Sheepdog - CNGA1 variant)	CNGA1	Deletion	0	AR	Clear
Progressive Retinal Atrophy (Discovered in the Swedish Vallhund)	MERTK	Insertion	0	AR	Clear
Progressive Retinal Atrophy 1 (Discovered in the Italian Greyhound)	Confidential	-	0	AR	Clear
Progressive Retinal Atrophy Type III	FAM161A	Insertion	0	AR	Clear
Protein Losing Nephropathy	NPHS1	G>A	0	AR	Clear
Pyruvate Dehydrogenase Phosphatase 1 Deficiency	PDP1	C>T	0	AR	Clear
Pyruvate Kinase Deficiency (Discovered in the Basenji)	PKLR	Deletion	0	AR	Clear
Pyruvate Kinase Deficiency (Discovered in the Beagle)	PKLR	G>A	0	AR	Clear
Pyruvate Kinase Deficiency (Discovered in the Pug)	PKLR	T>C	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Pyruvate Kinase Deficiency (Discovered in the West Highland White Terrier)	PKLR	Insertion	0	AR	Clear
QT Syndrome	KCNQ1	C>A	0	AD	Clear
Renal Cystadenocarcinoma and Nodular Dermatofibrosis	FLCN	A>G	0	AD	Clear
Rod-Cone Dysplasia 1	PDE6B	G>A	0	AR	Clear
Rod-Cone Dysplasia 1a	PDE6B	Insertion	0	AR	Clear
Rod-Cone Dysplasia 3	PDE6A	Deletion	0	AR	Clear
Sensorineural Deafness (Discovered in the Rottweiler)	LOXHD1	G>C	0	AR	Clear
Sensory Ataxic Neuropathy	tRNATyr	Deletion	0	MT	Clear
Sensory Neuropathy	FAM134B	Insertion	0	AR	Clear
Severe Combined Immunodeficiency (Discovered in Frisian Water Dogs)	RAG1	G>T	0	AR	Clear
Severe Combined Immunodeficiency (Discovered in Russell Terriers)	PRKDC	G>T	0	AR	Clear
Shaking Puppy Syndrome (Discovered in the Border Terrier)	Confidential	-	0	AR	Clear
Skeletal Dysplasia 2	COL11A2	G>C	0	AR	Clear
Spinocerebellar Ataxia (Late-Onset Ataxia)	CAPN1	G>A	0	AR	Clear
Spinocerebellar Ataxia with Myokymia and/or Seizures	KCNJ10	C>G	0	AR	Clear
Spondylocostal Dysostosis	HES7	Deletion	0	AR	Clear
Spongy Degeneration with Cerebellar Ataxia (Discovered in Belgian Malinois - SDCA1)	KCNJ10	T>C	0	AR	Clear
Spongy Degeneration with Cerebellar Ataxia (Discovered in Belgian Malinois - SDCA2)	ATP1B2	Insertion	0	AR	Clear
Stargardt Disease (Discovered in the Labrador Retriever)	ABCA4	Insertion	0	AR	Clear





Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
Startle Disease (Discovered in Irish Wolfhounds)	SLC6A5	G>T	0	AR	Clear
Startle Disease (Discovered in the Miniature American Shepherd)	Confidential	-	0	AR	Clear
Succinic Semialdehyde Dehydrogenase Deficiency (Discovered in the Saluki)	ALDH5A1	G>A	0	AR	Clear
Thrombopathia (Discovered in the Basset Hound)	RASGRP1	Deletion	0	AR	Clear
Thrombopathia (Discovered in the Eskimo Spitz)	RASGRP1	Insertion	0	AR	Clear
Trapped Neutrophil Syndrome	VPS13B	Deletion	0	AR	Clear
Van den Ende-Gupta Syndrome	SCARF2	Deletion	0	AR	Clear
von Willebrand's Disease, type 1	VWF	G>A	0	AD	Clear
von Willebrand's Disease, type 2	VWF	T>G	0	AR	Clear
von Willebrand's Disease, type 3 (Discovered in the Kooiker Hound)	VWF	G>A	0	AR	Clear
von Willebrand's Disease, type 3 (Discovered in the Scottish Terrier)	VWF	Deletion	0	AR	Clear
von Willebrand's Disease, type 3 (Discovered in the Shetland Sheepdog)	VWF	Deletion	0	AR	Clear
X-Linked Ectodermal Dysplasia	EDA	G>A	0	XR	Clear
X-Linked Hereditary Nephropathy (Discovered in the Navasota Dog)	COL4A5	Deletion	0	XR	Clear
X-Linked Hereditary Nephropathy (Discovered in the Samoyed)	COL4A5	G>T	0	XR	Clear
X-Linked Myotubular Myopathy	MTM1	C>A	0	XR	Clear
X-Linked Progressive Retinal Atrophy 1	RPGR	Deletion	0	XR	Clear
X-Linked Progressive Retinal Atrophy 2	RPGR	Deletion	0	XR	Clear
X-Linked Severe Combined Immunodeficiency (Discovered in the Basset Hound)	IL2RG	Deletion	0	XR	Clear



Breed: Lagotto Romagnolo  
Microchip number: 978101084931433  
Birth date: 2024-07-26

Registration number: FI39404/24  
Test date: 2024-12-14  
ID kit: DKGQLJC



Other health conditions tested

Genetic Condition	Gene	Risk Variant	Copies	Inheritance	Result
X-Linked Severe Combined Immunodeficiency (Discovered in the Cardigan Welsh Corgi)	IL2RG	Insertion	0	XR	Clear
X-Linked Tremors	PLP1	A>C	0	XR	Clear
Xanthinuria (Discovered in a mixed breed dog)	Confidential	-	0	AR	Clear
Xanthinuria (Discovered in the Cavalier King Charles Spaniel)	Confidential	-	0	AR	Clear
Xanthinuria (Discovered in the Toy Manchester Terrier)	Confidential	-	0	AR	Clear



## Glossary of genetic terms

### Test result definitions

**At Risk:** Based on the disorder's mode of inheritance, the dog inherited a number of genetic variant(s) which increases the dog's risk of being diagnosed with the associated disorder.

**Carrier:** The dog inherited one copy of a genetic variant when two copies are usually necessary to increase the dog's risk of being diagnosed with the associated disorder. While carriers are usually not at risk of clinical expression of the disorder, carriers of some complex variants may be associated with a low risk of developing the disorder.

**Clear:** The dog did not inherit the genetic variant(s) associated with the disorder and will not be at elevated risk of being diagnosed with the disorder due to this genotype. However, similar clinical signs could develop from different genetic or clinical causes.

**Inconclusive:** An inconclusive result indicates a confident call could not be made based on the data for that genetic variant. Health testing is performed in replicates, and on occasion the outcomes do not agree. This may occur due to an unusual sequence of DNA in the region tested, multiple cell genotypes present due to chimerism or acquired mutations, or due to quality of the DNA sample.

### Inheritance mode definitions

**Autosomal Recessive (AR):** For autosomal recessive disorders, dogs with two copies of the genetic variant are at risk of developing the associated disorder. Dogs with one copy of the variant are considered carriers and are usually not at risk of developing the disorder. However, carriers of some complex variants grouped in this category may be associated with a low risk of developing the disorder. Dogs with one or two copies may pass the disorder-associated variant to their puppies if bred.

**Autosomal Dominant (AD):** For autosomal dominant disorders, dogs with one or two copies of the genetic variant are at risk of developing the associated disorder. Inheriting two copies of the variant may increase the risk of development of the disorder or cause the condition to be more severe. These dogs may pass the disorder-associated variant to their puppies if bred.

**X-linked Recessive (XR):** For X-linked recessive disorders, the genetic variant is found on the X chromosome. Female dogs must inherit two copies of the variant to be at risk of developing the condition, whereas male dogs only need one copy to be at risk. Males and females with any copies of the variant may pass the disorder-associated variant to their puppies if bred.

**X-linked Dominant (XD):** For X-linked dominant disorders, the genetic variant is found on the X chromosome. Both male and female dogs with one copy of the variant are at risk of developing the disorder. Females inheriting two copies of the variant may be at higher risk or show a more severe form of the disorder than with one copy. Males and females with any copies of the variant may pass the disorder-associated variant to their puppies if bred.

**Mitochondrial (MT):** Unlike the two copies of genomic DNA held in the nucleus, there are thousands of mitochondria in each cell of the body, and each holds its own mitochondrial DNA (mtDNA). Mitochondria are called the "powerhouses" of the cell. For a dog to be at risk for a mitochondrial disorder, it must inherit a certain ratio of mtDNA with the associated variant compared to normal mtDNA. mtDNA is inherited only from the mother.